

Almost Coquaternion Structure

Constantin Udrişte,
Department of Mathematics IV,
Polytechnic Institute,
Bucharest, Romania

Abstract of the PhD Thesis in Mathematical Sciences, prepared under Prof. Dr. Gheorghe Th. Gheorghiu's guidance and defended at the Faculty of Mathematics and Mechanics of the Babeş-Bolyai University of Cluj, Romania, on September 28, 1971.

Preface

For some $(2m + 2)$ -dimensional differentiable manifolds one can define the notion of *almost complex structure* and for some $(4n + 4)$ -dimensional differentiable manifolds one can define the notion of *almost quaternion structure* by means of three almost complex structures satisfying certain conditions. These notions were first introduced by C. Ehresmann [7] in 1947 and then were studied by other geometricians [5], [6], [17], [20], [24], [25], [30], [48], [58], [59] etc, respectively [2], [22], [24], [26], [27], [28], [29], [57] etc.

For some $(2m + 1)$ -dimensional differentiable manifolds one can define the notion of *almost cocomplex structure* which may be considered as analogous to the almost complex structure (for even dimensional manifolds). This structure was first introduced in 1960 by S. Sasaki and called a (ϕ, ξ, η) -structure [34]. Later on, S. Sasaki and his students used the name *almost contact structure* for (ϕ, ξ, η) , as the structural group of the tangent bundle of a manifold which has a (ϕ, ξ, η) -structure may be reduced to $U(n) \times 1$ and conversely [5], [18], [19], [35], [41], [42], [58] etc. The same structure was studied in the papers of J. Bouzon [3], K. Ogiue and M. Okumura [31], K. Ogiue [30] etc, which named it an almost cocomplex structure.

The odd dimensional sphere S^{2m+1} is a typical example of differentiable manifold which admits an almost cocomplex structure (ϕ, ξ, η) [35]. We men-

tion that by means of the 1-form η which appears in the almost cocomplex structure of S^{2m+1} , Gh. Vranceanu [55] and C. Teleman [43] defined and studied on S^{2m+1} a nonholonomic manifold V_{2m+1}^{2m} whose Riemannian metric is that of a symmetric space V_{2m} . Taking this idea into account, we observed that on the almost cocomplex manifold a nonholonomic (holonomic) manifold of codimension one can be defined and studied by Pfaff's nonintegrable (complete integrable) equation $\eta = 0$.

The present paper defines and studies a structure for some $(4n + 3)$ -dimensional manifolds which is named almost coquaternion structure. This structure is composed of three almost cocomplex structures (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$, which satisfy some relations and may be considered as analogous to the almost quaternion structure for $(4n+4)$ -dimensional manifolds [49] – [53].

Independently, Y. Kuo [23] introduced in the same way the same structure calling it an *almost contact 3-structure*. Then, S. Sasaki [36], S. Tachibana and W. Yu [39], S. Tanno [40] etc investigated some properties of this structure using the denomination given by Kuo.

The sphere S^{4n+3} is a typical example of differentiable manifold which admits an almost coquaternion structure (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$ [49], [36]. Using the 1-forms η_a of the almost coquaternion structure of the sphere S^{4n+3} , C. Teleman [44] defined and studied on S^{4n+3} a nonholonomic manifold V_{4n+3}^{4n} whose Riemannian metric is the one of a symmetric space of E. Cartan [4]. Keeping in mind Teleman's idea, we observed that on an almost coquaternion manifold a nonholonomic (holonomic) manifold of codimension three can be defined and studied by nonintegrable (complete integrable) Pfaff's system $\eta_1 = 0, \eta_2 = 0, \eta_3 = 0$.

Our results are included in seven Chapters whose summary is given further on.

CONTENTS

Chapter I: Almost coquaternion structures

- §1 Algebraic preliminaries
- §2 Almost coquaternion structures
- §3 The structural group of the tangent bundle I
- §4 Associated Riemannian metric
- §5 Almost coquaternion metric structures
- §6 The structural group of the tangent bundle II

Chapter II: Almost coquaternion structures and theirs connection with almost quaternion structures

- §1 On the geometry of almost quaternion manifolds
- §2 The almost coquaternion structure of the manifold $\overline{M} \times R^3$
- §3 The almost coquaternion structure of the manifold $M \times R^3$
- §4 Almost quaternion contact manifolds and
their product with the almost coquaternion manifolds

Chapter III: Almost coquaternion affine connections

- §1 The bundle of coquaternion linear frames
- §2 Almost coquaternion affine connections
- §3 (ξ_a, η_a) -connections
- §4 Almost coquaternion metric connections
- §5 The holonomy group and the almost coquaternion structure

Chapter IV: Almost coquaternion hypersurfaces

- §1 Almost coquaternion hypersurfaces
- §2 Induced (ϕ_a, ξ_a, η_a) -connection
- §3 Almost coquaternion metric hypersurface
- §4 Imbedding of almost coquaternion Riemannian manifolds
- §5 A_m^0 -manifolds and η -Einstein manifolds

Chapter V: Fiberings of almost coquaternion manifolds

- §1 $Q_x M$ -invariant almost coquaternion structures
- §2 Fiberings of almost coquaternion manifolds
- §3 Fiberings of almost coquaternion Riemannian manifolds
- §4 The curvature form

Chapter VI: On ϕ -transformations

- §1 ϕ -transformations
- §2 Automorphisms
- §3 The almost coquaternion structure of
the $(4n + 3)$ -dimensional Lie groups

Chapter VII: Examples

- §1 The almost coquaternion structure of 3-dimensional orientable manifolds
- §2 The case of the sphere S^3
- §3 The case of the Tzitzeica hypersurface $xyzt = 1$
- §4 The sphere S^{4n+3} as an almost coquaternion manifold
- §5 The almost coquaternion structure of $T^1 \overline{M}$

References

CHAPTER I

We suppose that all the used differentiable manifolds and maps are of class C^∞ and we denote by $\mathcal{X}(M)$ the Lie algebra of all vector fields on the manifold M .

Let M be a $(4n + 3)$ -dimensional manifold.

1.1. Definition [49], [23]. *An almost coquaternion structure on M is an aggregate consisting of three almost cocomplex structures (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$, which satisfy*

$$\phi_a \circ \phi_b - \xi_a \otimes \eta_b = -\phi_b \cdot \phi_a + \xi_b \otimes \eta_a = \phi_c$$

$$\phi_a \xi_b = -\phi_b \xi_a = \xi_c$$

$$\eta_a \cdot \phi_b = -\eta_b \cdot \phi_a = \eta_c,$$

$$\eta_a(\xi_b) = \eta_b(\xi_a) = 0,$$

for any cyclic permutation $\{a, b, c\}$ of $\{1, 2, 3\}$.

In the paper [39] it is shown that the aggregate (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$, cannot be completed with a fourth almost cocomplex structure which, together with the three structures given above satisfy relations of the same type as those of Definition 1.1. On the other hand, we observe [49], [23] that in order to have an almost coquaternion structure on M it is sufficient to give only two almost cocomplex structures which satisfy certain relations, since the third structure results from the given structures. The more explicit form is [49]:

1.2. Theorem. *If a differentiable manifold M admits two almost cocomplex structures (ϕ_a, ξ_a, η_a) , $a = 1, 2$, which satisfy*

$$\phi_1 \circ \phi_2 + \phi_2 \circ \phi_1 = \xi_1 \otimes \eta_2 + \xi_2 \otimes \eta_1,$$

$$\eta_1(\xi_2) = 0,$$

then we have $\eta_2(\xi_1) = 0$, $\phi_1(\xi_2) = -\phi_2\xi_1$, $\eta_1 \circ \phi_2 = -\eta_2 \circ \phi_1$, and M admits a third almost cocomplex structure (ϕ_3, ξ_3, η_3) defined by

$$\phi_3 = \phi_1 \circ \phi_2 - \xi_1 \otimes \eta_2 = -\phi_2 \circ \phi_1 + \xi_2 \otimes \eta_1,$$

$$\xi_3 = \phi_1\xi_2 = -\phi_2\xi_1, \quad \eta_3 = \eta_1 \circ \phi_2 = -\eta_2 \circ \phi_1,$$

so that the relations of Definition 1.1 are valid.

1.3. Theorem. *If (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$, is an almost coquaternion structure on M , then $\forall (A_d^a) \in SO(3)$, $\forall \alpha : M \rightarrow R \setminus \{0\}$,*

$$\left(\bar{\phi}_d = \sum_a A_d^a \phi_a, \bar{\xi}_d = \frac{1}{\alpha} \sum_a A_d^a \xi_a, \bar{\eta}_d = \alpha \sum_a A_d^a \eta_a \right), \quad d = 1, 2, 3,$$

is again an almost coquaternion structure on M .

M is said to be an *almost coquaternion manifold*.

Let x be an arbitrary point of M . We denote the tangent space at x by $T_x M$ and its hypercomplexification $H \otimes_R T_x M$ by $T_x^H M$, where H is the corps of quaternion numbers and R is the real number field.

1.4. Theorem. *If by $D_{ax} M$, $a = 1, 2, 3$, we understand unidimensional distributions defined respectively by vector fields ξ_a , then the existence of an almost coquaternion structure on M is equivalent to the existence of the disjoint distributions $S_x^H M$, $D_{ax} M$ such that*

$$T_x^H M = S_x^H M \oplus \tau' S_x^H M \oplus \tau'' S_x^H M \oplus \tau''' S_x^H M \oplus D_{1x}^H M \oplus D_{2x}^H M \oplus D_{3x}^H M.$$

Thus, the existence of an almost coquaternion structure on M implies $\dim M = 3n + 3$.

Every almost coquaternion manifold M admits an *associated Riemannian metric* to the given almost coquaternion structure in the sense of

1.5. Theorem. *If M is an almost coquaternion manifold, then there exists a positive definite Riemannian metric g such that*

$$\eta_a(X) = g(\xi_a, X),$$

$$g(\phi_a X, \phi_a Y) = g(X, Y) - \eta_a(X) \eta_a(Y), \quad a = 1, 2, 3, \quad \forall X, Y \in \mathcal{X}(M).$$

The aggregate $(\phi_a, \xi_a, \eta_a, g)$, $a = 1, 2, 3$, is called an almost coquaternion metric structure and M is said to be an almost coquaternion Riemannian manifold.

An almost coquaternion metric structure can be described by means of 1-forms η_a and 2-forms $\Theta_a(X, Y) = g(\phi_a X, Y)$. Furthermore, we have

1.6. Theorem. *If $(\phi_a, \xi_a, \eta_a, g)$, $a = 1, 2, 3$, is an almost coquaternion metric structure, then, $\forall (A_d^a) \in SO(3)$, $\forall \alpha : M \rightarrow (0, \infty)$,*

$$\left(\sum_a A_d^a \phi_a, \frac{1}{\alpha} \sum_a A_d^a \xi_a, \alpha \sum_a A_d^a \eta_a, \alpha g + (\alpha^2 - \alpha) \sum_a \eta_a \otimes \eta_a \right), \quad d = 1, 2, 3,$$

is again an almost coquaternion metric structure on M .

Also, we find

1.7. Theorem. *If M is an almost coquaternion (Riemannian) manifold, then the structural group of its tangent bundle TM reduces to $GL(n, H) \times 1 \times 1 \times 1$ ($Sp(n) \times 1 \times 1 \times 1$). The converse is also true.*

Consequently, every almost coquaternion manifold M is orientable.

Some of the results presented in this Chapter have been published in [49].

CHAPTER II

Let $\overline{M}$ be a $4n$ -dimensional differentiable manifold which possesses an almost quaternion Hermitian structure (J_a, G) , $a = 1, 2, 3$, and M be a manifold which admits an almost coquaternion metric structure $(\phi_a, \xi_a, \eta_a, g)$, $a = 1, 2, 3$.

2.1. Theorem. *The product manifold $\overline{M} \times M$ possesses an almost coquaternion metric structure $(\phi'_a, \xi'_a, \eta'_a, g)$ defined by*

$$\phi'_a \begin{pmatrix} X \\ Y \end{pmatrix} = \begin{pmatrix} J_a X \\ \phi_a Y \end{pmatrix}, \quad \forall X \in T_x \overline{M}, \quad \forall Y \in T_y M, \quad (x, y) \in \overline{M} \times M,$$

$$\xi'_a = \begin{pmatrix} 0 \\ \xi_a \end{pmatrix}, \quad \eta'_a = (0, \eta_a), \quad g' = \begin{pmatrix} G & 0 \\ 0 & g \end{pmatrix}.$$

Taking into account that the aggregate $(0, \frac{d}{dt}, dt, 1)$ can be regarded as an almost cocomplex metric structure on the real line R , we have

2.2. Theorem. *If M is an almost coquaternion Riemannian manifold, then the product manifold $M \times R$ has an almost quaternion Hermitian structure*

$$J_a = \begin{pmatrix} \phi_a & \xi_a \\ -\eta_a & 0 \end{pmatrix}, \quad G = \begin{pmatrix} g & 0 \\ 0 & 1 \end{pmatrix}$$

induced by $(\phi_a, \xi_a, \eta_a, g)$ and $(0, \frac{d}{dt}, dt, 1)$.

As in [35], we obtain

2.3. Theorem. *The almost coquaternion structure determines the following tensor fields on M*

$$N_a^1(X, Y) = N_a(X, Y, \phi_a) + 2d\eta_a(X, Y)\xi_a, \quad \text{of type } (1, 2),$$

$$N_a^2(X, Y) = -2\eta_a(\phi_a X, Y) - 2d\eta_a(X, \phi_a Y)\xi_a, \quad \text{of type } (0, 2),$$

$$N_a^3(X) = -(L_{\xi_a} \phi_a)(X), \text{ of type } (1, 1),$$

$$N_a^4(X) = -(L_{\xi_a} \eta_a)(X), \text{ of type } (0, 1), \forall X, Y \in \mathcal{X}(M),$$

where $N_a(X, Y; \phi_a)$ is the Nijenhuis tensor of the endomorphism ϕ_a .

The tensor fields satisfy some relations which corresponding to those satisfied by the Nijenhuis tensors attached respectively to the three almost complex structures J_a on $M \times R$. As an application of these relations, we deduce

2.4. Theorem. N_1^1, N_2^1, N_3^1 vanish if any two of them vanish.

On the other hand, using the structure tensor T of the almost quaternion structure on $M \times R$ we deduce that the almost coquaternion structure determines the following tensor fields on M

$$T^1(X, Y) = \frac{2}{3} \sum_a N_a^1(X, Y), \quad \text{of type } (1, 2),$$

$$T^2(X, Y) = \frac{2}{3} \sum_a N_a^2(X, Y), \quad \text{of type } (0, 2),$$

$$T^3(X, Y) = \frac{2}{3} \sum_a N_a^3(X, Y), \quad \text{of type } (1, 1),$$

$$T^4(X, Y) = \frac{2}{3} \sum_a N_a^4(X, Y), \quad \text{of type } (0, 1), \forall X, Y \in \mathcal{X}(M).$$

Notice now, that T^1, T^2, T^3, T^4 satisfy some relations which correspond to those satisfied by the structure tensor T attached to the almost quaternion structure on $M \times R$. These imply

2.5. Theorem. If any one of T^1, T^2, T^3 , vanishes, then T^4 vanishes. If T^1 vanishes, then all the other tensors T^2, T^3, T^4 vanish.

2.6. Theorem. T^1 vanishes if and only if N_1^1, N_2^1, N_3^1 simultaneously vanish.

From the former discussions, it appears that the almost coquaternion structure (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$, on M whose tensor T^1 vanishes (identically) corresponds to the pseudo-quaternion structure J_a , $a = 1, 2, 3$, on $M \times R$. Hence it consists of three normal almost cocomplex structures.

2.7. Definition. The almost coquaternion structure whose tensor T^1 vanishes identically is called a pseudo-coquaternion structure and the manifold with such a structure a pseudo-coquaternion on manifold.

The induced almost quaternion structure on $M \times R$ may be integrable. So, we admit

2.8. Definition. *If the almost quaternion structure J_a on $M \times R$ is integrable, then we say that the almost coquaternion structure (ϕ_a, ξ_a, η_a) on M is normal and that M is a normal almost coquaternion manifold.*

A normal almost coquaternion structure is a pseudo-coquaternion structure. The converse is not generally true.

Finally, we observe that if N is a $(4m + 1)$ -dimensional manifold which possesses an almost quaternion contact metric structure $(\varphi_a, \xi, \eta, h)$, $a = 1, 2, 3$, [16], then we have

2.9. Theorem. *The product $M \times N$ of an almost coquaternion Riemannian manifold M and an almost quaternion contact Riemannian manifold N is an almost quaternion Hermitian manifold with the structure (J_a, G) , $a = 1, 2, 3$, defined by*

$$J_a \begin{pmatrix} X \\ Y \end{pmatrix} = \begin{pmatrix} \phi_a X + \eta(Y)\xi_a \\ \varphi_a Y - \eta_a(X)\xi \end{pmatrix}, \quad \forall X \in T_x M, \quad \forall Y \in T_y N,$$

$$G = \begin{pmatrix} g & 0 \\ 0 & h \end{pmatrix}.$$

The results obtained previously are included in [53].

CHAPTER III

First we show that the following theorem is valid.

3.1. Theorem. *An almost coquaternion (metric) structure on M is equivalent to the existence of a G -structure $P_G(M)$, where G is the real representation of $GL(n, H) \times 1 \times 1 \times 1$.*

This G -structure was called the *bundle of coquaternion linear frames (the bundle of coquaternion orthonormal frames)*.

3.2. Definition. A linear - or affine - connection of M is called an almost coquaternion (metric) connection if it is determined by a connection in the bundle of coquaternion linear frames (the bundle of coquaternion orthonormal frames).

We have

3.3. Theorem. *A linear connection ∇ of an almost coquaternion manifold M with almost coquaternion structure (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$, is an almost coquaternion connection if and only if $\phi_1, \phi_2, \xi_1 (\implies \phi_3, \xi_2, \xi_3, \eta_a)$ are parallel with respect to ∇ .*

3.4. Theorem. ∇ is an almost coquaternion affine connection if and only if the connection 1-forms ω_S^R of the coquaternion frame fields

$$X_S : X_s, X_{s'} = \phi_1 X_s, X_{s''} = \phi_2 X_s, X_{s'''} = \phi_3 X_s, X_{4n+a} = \xi_a,$$

$$S = s, s', s'', s''', 4n + a; s = 1, 2, \dots, n,$$

are given by the matrix

$$\omega = \left(\begin{array}{cccc|ccc} \omega_1^r & \omega_{s'}^r & \omega_{s''}^r & \omega_{s'''}^r & & & \\ -\omega_{s'}^r & \omega_s^r & -\omega_{s'''}^r & \omega_{s''}^r & & & \\ -\omega_{s''}^r & \omega_{s'''}^r & \omega_s^r & -\omega_{s'}^r & & & \\ -\omega_{s'''}^r & -\omega_{s''}^r & \omega_{s'}^r & \omega_s^r & & & \\ \hline & & & & 0 & 0 & 0 \\ & & & & 0 & 0 & 0 \\ & & & & 0 & 0 & 0 \end{array} \right).$$

In fact X_S constitute a system of $4n + 3$ independent congruences in $U \subset M$ and $\omega^R = ds^R$. Therefore, using the connection 1-forms we can build up the components γ_{TS}^R of the coquaternion affine connection ∇ with respect to the congruences X_S . Taking into account the definition of these components [54], [10-14] expressed in the invariant language [48], we find

$$\gamma_{TS}^R = ds^R(\nabla_{X_T} X_S) = \omega_S^R(X_T).$$

So, ∇ is a coquaternion affine connection if and only if its components with respect to the congruences X_S are given by $4n + 3$ matrices having the form of ω .

We also prove the following theorems.

3.5. Theorem. Every almost coquaternion manifold M admits an affine connection $\hat{\nabla}$ such that:

- 1) ξ_a and η_a are parallel with respect to $\hat{\nabla}$,
- 2) the torsion $\hat{T}$ of $\hat{\nabla}$ is given by

$$\hat{T}(X, Y) = 2 \sum_a d\eta(X, Y) \xi_a, \quad \forall X, Y,$$

where X and Y are vector fields.

3.6. Theorem. *A linear connection ∇ of an almost coquaternion Riemannian manifold M with almost coquaternion metric structure $(\phi_a, \xi_a, \eta_a, g)$ is an almost coquaternion metric connection if and only if $\phi_1, \phi_2, g(\Rightarrow \phi_3, \xi_a, \eta_a)$ are parallel with respect ∇ .*

3.7. Theorem. *∇ is an almost coquaternion metric connection if and only if the connection 1-forms ω_S^R of the coquaternion orthonormal frame fields X_S are given by the matrix ω with the additional conditions $\omega_S^R = -\omega_R^S$.*

Some results concerning holonomy and almost coquaternion (metric) structure on M are given at the end of this chapter.

These results are included in [51].

CHAPTER IV

Suppose $\overline{M}$ ($\dim \overline{M} = 4n + 4$) has an almost quaternion structure J_a , $a = 1, 2, 3$, and M is an orientable hypersurface imbedded in $\overline{M}$ by $i : M \rightarrow \overline{M}$. Let TM denote the tangent bundle of M and $T_r \overline{M}$ the restriction of the tangent bundle of $\overline{M}$ to M . We denote by i_* the differential of the imbedding i so that i_* is the mapping $i_* : TM \rightarrow T_r \overline{M}$. Let ζ be a pseudonormal vector field defined along M , i.e., $\zeta \in T_r \overline{M}$ and $\zeta \notin TM$. Then we can find a mapping $i_*^{-1} : T_r \overline{M} \rightarrow TM$ and a 1-form ω defined on M such that

$$i_*^{-1} \circ i_* = id, \quad i_* \circ i_*^{-1} = Id - \zeta \otimes \omega,$$

$$\omega \circ i_* = 0, \quad i_*^{-1} \zeta = 0, \quad \omega(\zeta) = 1, \quad \omega(J_a \zeta) = 0,$$

where id is the identity on TM and Id is the identity on $T_r \overline{M}$.

4.1. Theorem. *An orientable hypersurface M of an almost quaternion manifold $\overline{M}$ has a naturally induced almost coquaternion structure*

$$\phi_a = i_*^{-1} \circ J_a \circ i_*, \quad \xi_a = i_*^{-1} J_a \zeta, \quad \eta_a = -\omega \circ J_a \circ i_*, \quad a = 1, 2, 3.$$

Let D be an affine connection on $\overline{M}$. The affine connection of $\overline{M}$ induces an affine connection ∇ and three fundamental tensor fields: h of type (0,2) of type (1,1), m of type (0,1), on M in a natural manner [8], [37]. We have

4.2. Theorem. *Let D be an almost quaternion connection on $\overline{M}$, i.e.,*

$$(D_X J_a)(Y) = 0, \quad \forall X, Y \in \mathcal{X}(\overline{M}).$$

The connection ∇ on M is an almost coquaternion connection if and only if $h = 0$, $l = 0$, $m = 0$.

Suppose $\overline{M}$ has an Hermitian metric G . Then the induced metric g is given by

$$g(X, Y) = G(i_*X, i_*Y), \quad \forall X, Y \in T_x M.$$

$\overline{M}$ is orientable and we assume that $\overline{M}$ is oriented. Then we can choose a differentiable field ζ of unit normal vectors over M if and only if M is orientable. The field ζ satisfies the equations $G(\zeta, \zeta) = 1$, $G(\zeta, i_*X) = 0$, $\forall X \in T_x M$ and $\omega(J_a \zeta) = 0$, $a = 1, 2, 3$, together with its associated 1-form $\omega(X) = G(\zeta, X)$.

4.3. Theorem. *An orientable hypersurface M of an almost quaternion Hermitian manifold $\overline{M}$ has a naturally induced almost coquaternion metric structure*

$$\phi_a = i_*^{-1} \circ J_a \circ i_*, \quad \xi_a = i_*^{-1} J_a \zeta, \quad \eta_a = -\omega \circ J_a \circ i_*, \quad a = 1, 2, 3.$$

$$g(X, Y) = G(i_*X, i_*Y), \quad \forall X, Y \in T_x M.$$

Particularly, any hypersurfaces in R^{4n+4} which has one of the properties:

- 1) is simply connected and connected,
- 2) can be defined implicitly by $M : f(x) = c$, where the differential df is not zero at any point of M ,
- 3) is convex,

is orientable and hence it possesses an almost coquaternion metric structure. For example, the sphere S^{4n+3} is a compact manifold which has an almost coquaternion metric structure and the hypersurface of Tzitzeica $x^1 x^2 \dots x^{4n+4} = 1$ is a noncompact manifold which has an almost coquaternion metric structure.

If R is the real line, then we have

4.4. Theorem. *An almost coquaternion Riemannian manifold M can be imbedded in the almost quaternion Hermitian manifold $M \times R$ as a totally umbilical or geodesic hypersurface.*

The above mentioned ideas arose from [41] and were published in [50].

Let M be a m -dimensional manifold and Γ_{jk}^i , $i, j, k, l = 1, 2, \dots, m$, be a connection on M . We denote the Ricci tensor field attached to the given connection by Γ_{ij} .

4.5. Definition [15]. *If $\nabla_k(\det(\Gamma_{ij})) = 0$, then the manifold M is called an A_m^0 -manifold.*

4.6. Theorem. *If $\det(\Gamma_{ij}) \neq 0$, then M is an A_m^0 -manifold if and only if*

$$\Gamma^{ij} \nabla_k \Gamma_{ij} - \Gamma^{ij} \Gamma_{jk}^l \Gamma_{ei} + \Gamma_{lk}^l = 0, \quad \Gamma^{ij} \stackrel{def}{=} \frac{1}{\det(\Gamma_{ij})} \frac{\partial \det(\Gamma_{ij})}{\partial \Gamma_{ij}},$$

or

$$\Gamma^{ij}\nabla_k\Gamma_{ij} = 0,$$

if Γ_{ij} is a symmetrical tensor.

Using this Theorem we have:

4.7. Theorem. *Any η -Einstein manifold is an A_m^0 -manifold.*

CHAPTER V

Being given an almost coquaternion manifold M with structure (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$, we suppose that $\{\xi_a\}$ gives rise to a Lie group of transformations on M . Let us denote by G this 3-dimensional group.

5.1. Definition. *An almost coquaternion structure (ϕ_a, ξ_a, η_a) is called Q_xM -invariant if the distribution*

$$Q_xM = \{X | X \in T_xM, \eta_1(X) = 0, \eta_2(X) = 0, \eta_3(X) = 0\}$$

is invariant with respect to R_t , where R_t means the right translation in M by $t \in G$.

5.2. Theorem. *An almost coquaternion structure is Q_xM -invariant if and only if*

$$L_{\xi_1}\eta_a = C_{21}^a\eta_2 + C_{31}^a\eta_3,$$

$$L_{\xi_2}\eta_a = C_{12}^a\eta_1 + C_{32}^a\eta_3,$$

$$L_{\xi_3}\eta_a = C_{13}^a\eta_1 + C_{23}^a\eta_2,$$

where C_{bc}^a are the structure constants of the Lie algebra of G and L_{ξ_a} represents the Lie differentiation with respect to ξ_a .

5.3. Lemma. *If the structure (ϕ_a, ξ_a, η_a) is Q_xM -invariant, then the tensor field $\Gamma = \sum_a \xi_a \otimes \eta_a$ is invariant under the action of G . The converse is also true.*

As the tensor field of type (2,2)

$$P = \frac{1}{4}(id \otimes id + \sum_a \phi_a \otimes \phi_a + \Gamma \otimes \Gamma)$$

is invariant by G if and only if the field $\sum_a \phi_a \otimes \phi_a$ invariant by G , we admit

5.4. Definition. A $Q_x M$ -invariant almost coquaternion structure is said to be P -invariant if and only if $\sum_a \phi_a \otimes \phi_a$ is invariant under the action of G .

Let us suppose that M is an almost coquaternion Riemannian manifold with structure $(\phi_a, \xi_a, \eta_a, g)$, $a = 1, 2, 3$, and that $\{\xi_a\}$ determine a Lie group of motions with respect to g . In these conditions we get

5.5. Theorem. The group G is necessarily isomorphic to a unitary, semi-simple group, if $C_{23}^1 = m \neq 0$, or is isomorphic to an Abelian group, if $C_{23}^1 = 0$. If G is isomorphic to a unitary, semi-simple group, then its representative Riemannian space has a positive constant curvature.

We specify that in this case the structure (ϕ_a, ξ_a, η_a) is $Q_x M$ -invariant as

$$L_{\xi_a} \eta_a = 0, \quad L_{\xi_a} \eta_b = -L_{\xi_b} \eta_a = m \eta_c,$$

for any cyclic permutation $\{a, b, c\}$ of $\{1, 2, 3\}$. Evidently, the tensor field $\Gamma = \sum_a \xi_a \otimes \eta_a$ is invariant with respect to such a group G and we have

5.6. Lemma. If $\{\xi_a\}$ generates a unitary, semi-simple group of motions, with respect to g , such that

$$L_{\xi_a} \phi_a = 0, \quad L_{\xi_a} \phi_b = -L_{\xi_b} \phi_a = m \phi_c,$$

for any cyclic permutation $\{a, b, c\}$ of $\{1, 2, 3\}$, then the tensor fields

$$\begin{aligned} \sum_a \phi_a \otimes \phi_a, & \quad \text{of type } (2, 2), \\ P = \frac{1}{4} \left(id \otimes id + \sum_a \phi_a \otimes \phi_a + \Gamma \otimes \Gamma \right), & \quad \text{of type } (2, 2), \\ \phi_1 \otimes \phi_2 \otimes \phi_3, & \quad \text{of type } (3, 3), \\ \sum_a \xi_a \otimes \phi_a, & \quad \text{of type } (2, 1), \end{aligned}$$

are invariant under the action of G .

Let M be an almost coquaternion manifold with structure (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$. We admit

5.7. Definition. If $\{\xi_a\}$ gives rise to a Lie group of transformations on M and the distribution defined by $\{\xi_a\}$ is regular in Palais' sense [32], then the almost coquaternion structure (ϕ_a, ξ_a, η_a) is said to be strictly regular.

Suppose M is a connected compact manifold. We denote by M_G the quotient space and by $\pi : M \rightarrow M_G$ the canonical projection. It is known that [32] M_G is a connected compact manifold for which $\pi(U)(x^1, \dots, x^{4n})$ is a coordinate neighborhood.

5.8. Theorem. *If (ϕ_a, ξ_a, η_a) is a strictly regular and $Q_x M$ -invariant almost coquaternion structure on a connected compact manifold M , then*

- (i) *M is a principal G -bundle over M_G whose fiber is compact [32],*
- (ii) *$\Gamma = \sum_a \xi_a \otimes \eta_a$ is a connection on M , and*
- (iii) *$\eta = \{\eta_a\}$ is a connection 1-form on M .*

The Theorem 5.8 shows that we may consider principal bundles of type $M(M_G, G, \pi)$, where the total space M of possesses a $Q_x M$ -invariant almost coquaternion structure and G is the group generated by $\{\xi_a\}$.

5.9. Theorem. *If $M(M_G, G, \pi)$ is a principal bundle in which the total space M possesses a P -invariant structure (ϕ_a, ξ_a, η_a) then the tensor field*

$$P = \frac{1}{4} \left(id \otimes id + \sum_a \phi_a \otimes \phi_a + \Gamma \otimes \Gamma \right)$$

induces on M_G a global projector on the tensors of degree 2,

$$\mathcal{P}_p(X, Y) = d\pi(P_x)(X_x^*, Y_x^*),$$

$\forall p \in M_G, x \in M, p = \pi(x), X_x^*$ denotes the lift of $X \in \mathcal{X}(M_G)$ with respect to the connection η .

Suppose now that M possesses an almost coquaternion metric structure $(\phi_a, \xi_a, \eta_a, g)$, $a = 1, 2, 3$. Extending a result from [49] we get

5.10. Theorem. *If the connected compact manifold M has an almost coquaternion metric structure for which: (1) $\Theta_a = d\eta_a$, (2) the induced almost quaternion structure on $M \times R$ is a pseudo-quaternion structure, (3) the distribution defined by $\{\xi_a\}$ is regular in Palais' sense [32], then*

(i) *M is a principal bundle whose fiber is a sphere S^3 or a real projective space $RP^3 = S^3/\{id., -id\}$.*

(ii) *$\Gamma = \sum_a \xi_a \otimes \eta_a$ is a connection on M , and*

(iii) *$\eta = \{\eta_a\}$ is a connection 1-form on M .*

As g is a G -invariant Riemannian metric on M , we can define a Riemannian metric on M_g by

$$h(X, Y) = g(X^*, Y^*), \quad X, Y \in \mathcal{X}(M_G),$$

called the *induced Riemannian metric*.

Next we suppose that the hypotheses of Theorem 5.10 are satisfied and hence Lemma 5.6 is valid. Therefore all the tensor fields mentioned in lemma 5.6 induce tensor fields on M_G whose properties are included in some theorems.

The Theorems stated in this part of the paper hold good for any principal bundle $M(M_G, G, \pi)$ in which the total space M possesses an almost coquaternion metric structure and D is a unitary semi-simple group of motion satisfying Lemma 5.6.

Finally, we make some observations on the structure equation of the fibering in Theorem 5.10.

CHAPTER VI

Let M be an almost cocomplex manifold with structure (ϕ, ξ, η) and f a transformation of M .

6.1. Definition. *A transformation f of M which leaves ϕ invariant, i.e.,*

$$f_* \circ \phi = \phi \circ f_*$$

is called a ϕ -transformation [35].

6.2. Lemma. *If f is a ϕ -transformation of M , then there exists a function $\varepsilon : M \rightarrow R$ such that*

$$f_* \xi = \varepsilon \xi, \quad f^* \eta = \varepsilon \circ f \eta.$$

Consider a subset of infinitesimal ϕ -transformations X such that there is a system of differential equations

$$(*) \quad \eta_k \xi^i \frac{\partial^2 X^k}{\partial x^i \partial x^j} = h_{kj}^i(x) \frac{\partial X^k}{\partial x^i} + h_{kj}(x) X^k + h_j(x)$$

defined in a neighborhood of $x \in M$ and satisfied by components X^i of X ; denote by $\mathcal{T}_0(M; \phi)$ the corresponding subset of ϕ -transformations in the set of all ϕ -transformations over M .

Sasaki's papers [35] suggested us to use some results of [1], [32] in order to prove the following

6.3. Theorem. *If M is a compact manifold, then $\mathcal{T}_0(M; \phi)$ is a Lie group of transformations.*

Let M be an almost coquaternion manifold with structure tensors (ϕ_a, ξ_a, η_a) , $a = 1, 2, 3$.

6.4. Definition. A transformation f on M which leaves ϕ_1 and ϕ_2 invariant, i.e.,

$$f_* \circ \phi_a = \phi_a \circ f_*, \quad a = 1, 2,$$

is called a (ϕ_1, ϕ_2) -transformation.

6.5. Lemma. If f is a (ϕ_1, ϕ_2) -transformation of M , then

$$f_* \circ \phi_3 = \phi_3 \circ f_*$$

and there exists a function $\varepsilon : M \rightarrow R$ such that

$$f_* \xi_a = \varepsilon \xi_a, \quad f^* \eta_a = \varepsilon \circ f \eta_a, \quad a = 1, 2, 3.$$

Let $\mathcal{T}_0(M; \phi_1, \phi_2)$ be the subset of all (ϕ_1, ϕ_2) -transformations which verify the relation (*). We have

6.6. Corollary. If M is a compact almost coquaternion manifold, then $\mathcal{T}_0(M; \phi_1, \phi_2)$ is a Lie group of transformations.

Let M be an almost cocomplex manifold.

6.7. Definition. A transformation f of M which leaves the structure (ϕ, ξ, η) invariant, i.e.,

$$f_* \circ \phi = \phi \circ f_*, \quad f_* \xi = \xi \quad (\Rightarrow f^* \eta = \eta)$$

is called an automorphism of M [35].

It results: if M is a compact manifold, then the set of all automorphisms of M is a Lie group [35].

Let M be an almost coquaternion manifold.

6.8. Definition. A transformation f of M which leaves the structure (ϕ_a, ξ_a, η_a) invariant, i.e.,

$$f_* \circ \phi_1 = \phi_1 \circ f_*, \quad f_* \circ \phi_2 = \phi_2 \circ f_*, \quad f_* \xi_1 = \xi_1,$$

$$(\Rightarrow f_* \circ \phi_3 = \phi_3 \circ f_*, \quad f_* \xi_2 = \xi_2, \quad f_* \xi_3 = \xi_3, \quad f^* \eta_a = \eta_a, \quad a = 1, 2, 3),$$

is called an automorphism of M .

It follows that the set of all automorphisms of a compact almost coquaternion manifold M is a Lie group.

Finally, we prove

6.9. Theorem. *Every $(4n + 3)$ -dimensional Lie group admits a left invariant almost coquaternion structure.*

The results obtained in this chapter have been included in [52].

CHAPTER VII

Let M be a 3-dimensional manifold. We have

7.1. Theorem. *M has an almost coquaternion (metric) structure if and only if it is parallelizable.*

The hypothesis that M is parallelizable is equivalent to the fact that it possesses three vector fields ξ_a which are linearly independent at every point of M . Let η_a be the dual 1-forms. Define three linear endomorphisms ϕ_a by

$$\phi_1 = \xi_3 \otimes \eta_2 - \xi_2 \otimes \eta_3, \phi_2 = \xi_1 \otimes \eta_3 - \xi_3 \otimes \eta_1, \phi_3 = \xi_2 \otimes \eta_1 - \xi_1 \otimes \eta_2$$

and a metric tensor g by

$$g = \eta_1 \otimes \eta_1 + \eta_2 \otimes \eta_2 + \eta_3 \otimes \eta_3.$$

$(\phi_a, \xi_a, \eta_a, g)$, $a = 1, 2, 3$, defines an almost coquaternion metric structure on M .

As any orientable 3-dimensional manifold is parallelizable [38], we have

7.2. Theorem. *Every 3-dimensional orientable manifold has an almost coquaternion metric structure.*

The above presented results permit us to give explicitly almost coquaternion metric structures on the sphere $S^3 : x^2 + y^2 + z^2 + t^2 = 1$ and on the Tzitzeica hypersurface $xyzt = 1$.

Then it is shown that the sphere S^{4n+3} possesses a normal coquaternion metric structure $(\phi_a, \xi_a, \eta_a, g)$, $a = 1, 2, 3$, for which $\Theta_a = d\eta_a$ and the distribution defined by $\{\xi_a\}$ is regular in Palais' sense. So the sphere S^{4n+3} is an example of our results obtained in Chapter V. The quotient space of this principal bundle is the quaternionic projective space [2], [54].

Finally, we construct some almost coquaternion structures on some hypersurfaces of the tangent bundle of an almost quaternion (Hermitian) manifold.

The author expresses his heartiest thanks to Professor Gheorghe Th. Gheorghiu for his kind remarks and valuable advices in preparing the PhD Thesis.

References

- [1] Bochner, S., *Tensor fields with finite bases*, Ann. of Math. (2), 53 (1951), pp. 400-411.
- [2] Bonan, E., *G-structures de type quaternionien*, Thèse, Paris (1967).
- [3] Bouzon, J., *Structures presque-cocomplexes*, Thèse, Paris (1965).
- [4] Cartan, E., *Sur certaines formes Riemanniennes remarquables des géométries à groupe fondamental simple*, Annales de l'Ecole Normale Sup., 44 (1927), pp. 348.
- [5] Chern, S., *The geometry of G-structures*, Bull. Amer. Math. Soc., 72 (1966), pp. 167-219.
- [6] Clark, R.S. and Bruckheimer, M., *Tensor structures on a differentiable manifold*, Ann. Mat. Pura Appl. (4), 54 (1961), pp. 123-141.
- [7] Ehresmann, C., *Sur la théorie des espaces fibrés*, Colloque de Topologie Algébrique, C.N.R.S., Paris (1947), pp. 3.
- [8] Eisenhart, L.P., *Non-Riemannian Geometry*, Amer. Math. Soc., New York (1958).
- [9] Gheorghiu, Th. Gh., *Hipersuprafețe Țițeica*, Lucările științifice ale Institutului Pedagogic Timișoara (1959).
- [10] Gheorghiu, Th. Gh., *Sur un calcul tensoriel mixte*, Atti della Acc. Sci. di Bologna, Rendiconti, Serie XII, t. III (1965-1966).
- [11] Gheorghiu, Th. Gh., *Sur les courbes géodésique d'un calcul absolute mixte*, Analele St. Univ. Iași, Matematică, t. XI-B (1965), pp. 273-281.
- [12] Gheorghiu, Th. Gh., *Connexions projectives dans un calcul absolute mixte*, Rend. Sem. Mat. Messina, t. XI (1966-1967), pp. 21-32.
- [13] Gheorghiu, Th. Gh., *Des coordonnées normales dans un calcul absolut mixte relativement à un espace Riemannien*, Rev. Roum. Math. Pures Appl. 13 (1968), pp. 451-468.
- [14] Gheorghiu, Th. Gh., *Des coordonnées normales mixtes dans un espace à connexion affine*, Rend. Sem. Mat. Fis. di Milano XXXVIII (1968).

- [15] Gheorghiu, Th. Gh., *O clasă particulară de spații cu conexiune afină*, St. Cerc. Mat. t. 21, 8 (1969), pp. 1157-1168.
- [16] Hashimoto, S., *On differentiable manifolds with almost quaternion contact structure*, Tensor (N.S.) 15 (1964), pp. 258-268.
- [17] Ianuș, S., și Udriște, C., *Asupra spațiului fibrat tangent al unei varietăți diferențiabile*, St. Cerc. Mat. t. 22, 4 (1970), pp. 599-611.
- [18] Ianuș, S., *Quelques remarques sur l'espace fibré tangent d'une variété cosymplectique*, An. St. Univ. Iași, Matematică (1971).
- [19] Ianuș, S., *Some almost product structure on manifolds with linear connections*, Kodai Math. Sem. Rep. (1971).
- [20] Kobayashi, S. and Nomizu, K., *Foundations of differential geometry*, vol. I, II, Interscience, New York (1963) and (1969).
- [21] Koszul, J.L., *Lectures on fibre bundles and differential geometry*, Tata Inst. Fund. Research, Bombay (1960).
- [22] Kraines, V.Y., *Topology of quaternionic manifolds*, Bull. Amer. Math. Soc. 71 (1965), pp. 526-527; same title, Trans. Amer. Math. Soc. 122 (1966), pp. 357-367.
- [23] Kuo, Y.Y., *On almost contact 3-structure*, Tôhoku Math J. 22 (1970), pp. 325-332.
- [24] Libermann, P., *Sur les structures presque complexes et autres structures infinitésimales régulières*, Bull. Soc. Math. France 83 (1955).
- [25] Lichnerowicz, A., *Théorie globale des connexions et des groupes d'holonomie*, Rome (1955).
- [26] Martinelli, E., *Varieta a struttura quaternionale generalizzata*, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Nat. 26 (1959), pp. 353-362.
- [27] Martinelli, E., *Modello metrico reale dello spazio proiettivo quaternionale*, An. Mat. Pura Appl. 49 (1960), pp. 78-89.
- [28] Obata, M., *Affine connections on manifolds with almost complex, quaternion or Hermitian structure*, Jap. J. Math. 26 (1956), pp. 43-77.

- [29] Obata, M., *Hermitian manifolds with quaternion structure*, Tôhoku Math. J. 10 (1958).
- [30] Ogiue, K., *G-structure defined by tensor fields*, Kôdai Math. Sem. Rep. 20 (1968), pp. 54-76.
- [31] Ogiue, K. and Okumura, M., *On cocomplex structures*, Kôdai Math. Sem. Rep. 19 (1967), pp. 507-512.
- [32] Palais, R.S., *A global formulation of the Lie theory of transformations groups*, Mem. Amer. Math. Soc. 22 (1957).
- [33] Papuc, I.D. și Albu, C.A., *Elemente de geometrie diferențială globală*, Tipografia Univ. Timișoara (1970).
- [34] Sasaki, S., *On differentiable manifolds with certain structures which are closely related to almost contact structures*, Tôhoku Math. J. 12 (1960), pp. 459-476; II (with Hatakeyama, Y.) 13 (1961), pp. 281-294.
- [35] Sasaki, S., *Almost contact manifolds*, Math. Inst. Tôhoku Univ. Part I (1965), Part II (1967), Part III (1968).
- [36] Sasaki, S., *On spherical space forms with normal contact metric 3-structure*, manuscripts.
- [37] Schouten, J.A., *Ricci-Calculus*, Berlin (1954).
- [38] Stiefel, E., *Richtungsfelder und fernparallelisms in n-dimensionalen Mannigfaltigkeiten*, Comment. Math. Helv. 8 (1936), pp. 305-353.
- [39] Tachibana, S. and Yu, W.N., *On a Riemannian space admitting more than one Sasakian structures*, Tôhoku Math. J. 22 (1970), pp. 535-540.
- [40] Tanno, S., *On the isometry groups of Sasakian manifolds*, J. Math. Soc. Japan, 22 (1970), pp. 579-590.
- [41] Tashiro, Y., *On contact structure of hypersurfaces in complex manifolds I*, Tôhoku Math. J. 15 (1963), pp. 62-78.
- [42] Tashiro, Y., *On contact structure of tangent sphere bundles*, Tôhoku Math. J. 21 (1969), pp. 117-143.

- [43] Teleman, C., *Asupra grupurilor de mișcare maxime ale spațiilor riemanniene V_n* , St. Cerc. Mat. 5(1954), pp. 143-171.
- [44] Teleman, C., *Sur certaines espaces symétriques*, Bulletin Scientifique, Acad. R.P.R. t.7, 4 (1955), pp. 977-1002.
- [45] Țițeica, G., *Sur quelques propriétés affines*, Bull. Math. Phys. de l'Ec. Pol., București, t. 6(1936), pp. 8-10.
- [46] Țițeica, G., *Opere*, vol. I, București (1941).
- [47] Udriște, C., *Introducerea unor invarianti diferențiali intrinseci într-o varietate riemanniană*, St. Cerc. Mat. 21 (1969), pp. 509-514.
- [48] Udriște, C., *Congruences on the tangent bundle of a differentiable manifold*, Rev. Roum. Math. Pures Appl., t. 15 (1970), pp. 1079-1096.
- [49] Udriște, C., *Structures presque coquaterniennes*, Bull. Math. de la Soc. Sci. math. de la R.S.R., t. 13 (61), 4(1969), pp. 487-507. O parte din această lucrare a fost prezentată la "A' IV-lea congres al matematicienilor de expresie latină", București, 17-24 septembrie 1969.
- [50] Udriște, C., *Almost coquaternion hypersurfaces*, Rev. Roum. Math. Pures Appl., t. 15, 9(1970), pp. 1545-1553.
- [51] Udriște, C., *Coquaternion structures and almost coquaternion structures*, Rev. Roum. Math. Pures Appl. (sub tipar).
- [52] Udriște, C., *On ϕ -transformations*, An. St. Univ. Iași, Matematică (1971).
- [53] Udriște, C., *On almost coquaternion structures*, Studia Univ. Babeș-Bolyai, series Math. Phys., Cluj (1971).
- [54] Vrănceanu, Gh., *Lecții de geometrie diferențială*, vol. I, II (1964).
- [55] Vrănceanu, Gh., *Asupra unei clase de spații riemanniene omogene*, St. Cerc. Mat., t. 5, 1-2 (1954), pp. 173-223.
- [56] Vrănceanu, Gh., *Varieta differenziabili anolonome*, Rend. Sem. Mat. Messina, t. 10 (1965-1966), pp. 139-146.

- [57] Wakakuwo, H., *On Riemannian manifolds with homogeneous holonomy groups $Sp(n)$* , Tôhoku Math. J. 10 (1958).
- [58] Yano, K., *On a structure defined by a tensor field f of type $(1, 1)$ satisfying $f^3 + f = 0$* , Tensor, 14 (1965), pp. 99-109.
- [59] Yano, K., *Differential geometry on complex and almost complex spaces*, Pergamon Press (1965).